PRODUCT INFORMATION



1-hydroxy Vitamin D2

Item No. 12077

CAS Registry No.: 54573-75-0

Formal Name: 4-methylene-5Z-[2E-[(1R,3aS)-octahydro-7aR-

> methyl-1R-[1R,4,5-trimethyl-2E-hexen-1-yl]-4Hinden-4-ylidene]ethylidene]-1,3S-cyclohexanediol

Synonyms: Doxercalciferol, Hectorol, 1-Hydroxyergocalciferol,


TSA 840

MF: $C_{28}H_{44}O_{2}$ FW: 412.7 **Purity:** ≥98%

 λ_{max} : 211, 265 nm UV/Vis.: A crystalline solid Supplied as:

-20°C Storage: ≥4 years Stability:

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

1-hydroxy Vitamin D₂ is supplied as a crystalline solid. A stock solution may be made by dissolving the 1-hydroxy vitamin \bar{D}_2 in the solvent of choice, which should be purged with an inert gas. 1-hydroxy Vitamin D₂ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of 1-hydroxy vitamin D₂ in these solvents is approximately 20 mg/ml.

1-hydroxy Vitamin D_2 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 1-hydroxy vitamin D_2 should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. 1-hydroxy Vitamin \overline{D}_2 has a solubility of approximately 0.3 mg/ml in a 1:2 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Vitamin D aids in the absorption of calcium and has central roles in bone formation and maintenance, hypertension, cancer, and immunity. 1,2 1-hydroxy Vitamin D_2 is a synthetic prodrug of the active 1,25-dihydroxy vitamin D_2 (Item No. 11174), as well as of other active metabolites.³ It is used to suppress the synthesis and secretion of parathyroid hormone in secondary hyperparathyroidism, an action which requires vitamin D receptor.⁴⁻⁶ 1-hydroxy Vitamin D₂ is effective in stimulating bone growth and, in combination therapy with angiotensin receptor blockers, ameliorates albuminuria and glomerulosclerosis associated with diabetic nephropathy.^{7,8} It also inhibits the growth of certain blastomas and prostate cancer cells.^{9,10}

References

- 1. Holick, M.F. N. Engl. J. Med. 357(3), 266-281 (2007).
- 2. Peterlik, M., Boonen, S., Cross, H.S., et al. Int. J. Environ. Res. Public Health 6, 2585-607 (2009).
- 3. Upton, R.A., Knutson, J.C., Bishop, C.W., et al. Nephrol. Dial. Transplant. 18, 750-758 (2003).
- 4. Frazao, J.M., Chesney, R.W., Coburn, J.W., et al. Nephrol. Dial. Transplant. 13, 68-72 (1998).
- 5. Brown, A.J., Dusso, A.S., and Slatopolsky, E. Nephrol. Dial. Transplant. 17, 10-19 (2002).
- 6. Ritter, C.S. and Brown, A.J. J. Mol. Endocrinol. 46(2), 63-66 (2011).
- 7. Nguyen-Yamamoto, L., Bolivar, I., Strugnell, S.A., et al. J. Am. Soc. Nephrol. 21, 1713-1723 (2010).
- Zhang, Y., Deb, D.K., Kong, J., et al. Am. J. Physiol. Renal Physiol. 297, F791-F801 (2009).
- 9. van Ginkel, P.R., Yang, W., Marcet, M.M., et al. J. Neurooncol. 85, 255-262 (2007).
- 10. Attia, S., Eickhoff, J., Wilding, G., et al. Clin. Cancer Res. 14(8), 2437-2443 (2008).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 10/07/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM