PRODUCT INFORMATION




COX-1 (ovine) Electrophoresis Standard

Item No. 360100

Overview and Properties

Contents:	This vial contains 5 μg purified COX-1.
Synonyms:	Cyclooxygenase 1, PGHS-1, Prostaglandin Endoperoxide Synthase 1, Prostaglandin
	G/H Synthase 1, Prostaglandin H2 Synthase 1
Form:	Liquid
Storage:	-80°C (as supplied)
Stability:	≥1 year
Storage Buffer:	20 mM Tris-HCl buffer, pH 8.0
Purity:	≥95%
Applications:	This enzyme has been denatured and has no catalytic activity. It can be used as a
	standard for Western blots (WB) and electrophoresis.

Image



WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/01/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Cyclooxygenase 1 (COX-1) is a bifunctional enzyme that exhibits both COX and peroxidase activities.^{1,2} It is composed of an N-terminal signal peptide, an EGF-like domain, a membrane binding domain, a catalytic domain, and a C-terminal tail.³ COX-1 is constitutively expressed in the gastrointestinal tract, kidney, spleen, liver, and lung and localizes to the endoplasmic reticulum.^{4,5} The COX component converts arachidonic acid (Item Nos. 90010 | 90010.1 | 10006607) to a hydroperoxyl endoperoxide prostaglandin G₂ (PGG₂; Item No. 17010) and the peroxidase component reduces the endoperoxide to the corresponding alcohol PGH₂ (Item No. 17020), the precursor of PGs, thromboxanes, and prostacyclins.^{1,2} COX-1 is the target of many non-steroidal anti-inflammatory drugs (NSAIDs) and is responsible for the undesirable gastrointestinal and renal side effects, such as ulcer formation and reductions in the glomerular filtration rate, respectively.^{6,7} This product is intended to be used as a standard for Western blot (WB) and electrophoresis. This enzyme has been carefully purified to exclude all unrelated proteins and may not be catalytically active.

References

- 1. Nugteren, D.H. and Hazelhof, E. Isolation and properties of intermediates in prostaglandin biosynthesis. *Biochim. Biophys. Acta* **326(3)**, 448-461 (1973).
- 2. Hamberg, M. and Samuelsson, B. Detection and isolation of an endoperoxide intermediate in prostaglandin biosynthesis. *Proc. Natl. Acad. Sci. USA* **70(3)**, 899-903 (1973).
- 3. Smith, W.L. and DeWitt, D.L. Prostaglandin endoperoxide H synthases-1 and -2. Adv. Immunol. 62, 167-215 (1995).
- 4. Seibert, K., Zhang, Y., Leahy, K., *et al.* Pharmacological and biochemical demonstration of the role of cyclooxygenase 2 in inflammation and pain. *Proc. Natl. Acad. Sci. USA* **91(25)**, 12013-12017 (1994).
- 5. Morita, I., Schindler, M., Regier, M.K., *et al.* Different intracellular locations for prostaglandin endoperoxide H synthase-1 and -2. J. Biol. Chem. **270(18)**, 10902-10908 (1995).
- 6. Gierse, J.K., Hauser, S.D., Creely, D.P., *et al.* Expression and selective inhibition of the constitutive and inducible forms of human cyclo-oxygenase. *Biochem. J.* **305(Pt 2)**, 379-484 (1995).
- 7. Frölich, J.C. A classification of NSAIDs according to the relative inhibition of cyclooxygenase isoenzymes. *Trends Pharmacol. Sci.* **18(1)**, 30-34 (1997).

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM